

Wnt/Frizzled Signaling Controls *C. elegans* Gastrulation by Activating Actomyosin Contractility

Jen-Yi Lee,^{1,4,5} Daniel J. Marston,^{1,4} Timothy Walston,² Jeff Hardin,^{2,3} Ari Halberstadt,¹ and Bob Goldstein^{1,*}

¹Department of Biology

University of North Carolina, Chapel Hill
Chapel Hill, North Carolina 27599-3280

²Laboratory of Genetics

³Program in Cellular and Molecular Biology
and Department of Zoology

University of Wisconsin-Madison
Madison, Wisconsin 53706

Summary

Background: Embryonic patterning mechanisms regulate the cytoskeletal machinery that drives morphogenesis, but there are few cases where links between patterning mechanisms and morphogenesis are well understood. We have used a combination of genetics, in vivo imaging, and cell manipulations to identify such links in *C. elegans* gastrulation. Gastrulation in *C. elegans* begins with the internalization of endodermal precursor cells in a process that depends on apical constriction of ingressing cells.

Results: We show that ingressation of the endodermal precursor cells is regulated by pathways, including a Wnt-Frizzled signaling pathway, that specify endodermal cell fate. We find that Wnt signaling has a role in gastrulation in addition to its earlier roles in regulating endodermal cell fate and cell-cycle timing. In the absence of Wnt signaling, endodermal precursor cells polarize and enrich myosin II apically but fail to contract their apical surfaces. We show that a regulatory myosin light chain normally becomes phosphorylated on the apical side of ingressing cells at a conserved site that can lead to myosin-filament formation and contraction of actomyosin networks and that this phosphorylation depends on Wnt signaling.

Conclusions: We conclude that Wnt signaling regulates *C. elegans* gastrulation through regulatory myosin light-chain phosphorylation, which results in the contraction of the apical surface of ingressing cells. These findings forge new links between cell-fate specification and morphogenesis, and they represent a novel mechanism by which Wnt signaling can regulate morphogenesis.

Introduction

The morphogenetic events that shape embryonic development rely on the movements and shape changes of individual cells. Because the cellular cytoarchitecture provides the driving forces for these cellular events,

one of the keys to understanding the molecular basis of morphogenetic movements is determining how well-studied developmental pathways specifying cell fate lead to modulation of the cytoskeleton in individual cells in ways that can produce forces capable of moving cells or deforming tissues. Toward this goal, there has been identification of many genes that function upstream of morphogenetic movements, including many essential for cell-fate specification and extracellular signaling, and there is some understanding of the cytoskeletal mechanics that drive these movements. However, there has been more limited progress in tying these two ends together to provide a coherent thread from cell fate and signaling molecules to the cytoskeletal dynamics responsible for morphogenesis [1, 2].

One of the earliest morphogenetic events in animal development is gastrulation, the process by which the embryo reorganizes itself into three germ layers. Gastrulation in *C. elegans* begins at the 26-cell stage when the two endodermal founder cells, Ea and Ep, begin to migrate from the outer, ventral surface of the embryo to the embryonic interior [3] (Figure 1). The mechanisms that specify endodermal fate in these cells are well studied. SKN-1, an endomesodermal determinant, is segregated to two cells (P₂ and EMS) at the four-cell stage. In P₂, SKN-1 activity is repressed in the P₂ cell, whereas SKN-1 activity persists in EMS to promote endomesodermal fate [4]. A Wnt interaction at the four-cell stage then specifies endodermal fate on one side of the EMS cell. Differential regulation of transcription factor activity in this cell's two daughters results in a single endodermal precursor cell (E) at the eight-cell stage [5].

After E is born, its daughter cells (Ea and Ep) ingress during a cell cycle that is extended by the introduction of a gap phase [6]. The space left on the ventral side of the embryo is filled by neighboring cells, a total of six cells from the MS, AB, and P₄ lineages [7]. After ingressation, Ea and Ep divide (Figure 2A). For simplicity, we use the term “gastrulation” here solely to refer to Ea-Ep ingressation, the internalization of the endoderm. Gastrulation in *C. elegans* continues later with the internalization of other cells including mesoderm and germline progenitors [8].

Recent work has begun to shed light on the mechanisms required for *C. elegans* gastrulation [9]. One of the driving forces for Ea-Ep ingressation is apical constriction, which is likely powered by an actomyosin contraction [7]. Consistent with this finding, NMY-2, a non-muscle myosin II heavy chain, accumulates at the apical (ventral) surfaces of Ea and Ep, and this polarized accumulation requires the activity of the PAR proteins [8, 10]. The PAR proteins were first identified as essential for polarity in the 1-cell *C. elegans* embryo [11] and have since been found to be required for polarity in organisms as diverse as *Drosophila* and humans [12]. In gastrulation-stage embryos, the PAR proteins have polarized distributions that are established by cell-cell contacts, with PAR-3, PAR-6, and PKC-3 localized apically, at

*Correspondence: bobg@unc.edu

⁴These authors contributed equally to this work.

⁵Present address: Department of Molecular and Cell Biology, Mail Code #3200, University of California, Berkeley, California 94720-3200




Figure 1. Three-Dimensional Illustrations of Embryos prior to and during Gastrulation

Ea and Ep (green) become completely enveloped by neighboring cells as gastrulation proceeds. These 3D renderings are based on tracings of optical sections of 24-cell (A) and 28-cell (B) embryos that were stained with labeled phalloidin to mark the cell cortex in each cell. Six cells extend into the gap vacated by the ingressing Ea and Ep cells. Two of these—one granddaughter of MS, left, and the P₄ cell, right—are shown here in opaque blue. Illustrations by Janet Iwasa (janet@onemicron.com).

contact-free areas, and PAR-1 and PAR-2 localized basolaterally where cells contact neighboring cells [8, 10]. Depleting the embryo of specific PAR proteins just before gastrulation compromises gastrulation movements. The timing of ingression is delayed significantly compared to that in wild-type cells, but the E lineage cells still internalize [10]. This suggests that there must be additional and essential gastrulation regulators that remain to be identified.

The Wnt pathway has been implicated in cell-fate specification, cell polarization, and morphogenesis across the animal kingdom [13]. Wnt ligands or their Frizzled (Fz) receptors, or both, are required in many processes including establishment of *Drosophila* segment polarity [14, 15], zebrafish and *Xenopus* gastrulation [16–18], and *Xenopus* neural tube closure [19]. Wnt and Fz are known to act through various signal transduction pathways, generally categorized as canonical and noncanonical pathways. Canonical signaling results in translocation of β -catenin from the cytoplasm to the nucleus, where it participates in the transcriptional activation of downstream targets. In *C. elegans*, *mom-2*, the Wnt gene that functions during the four-cell stage to specify endoderm, acts in a variant of the canonical manner by activating WRM-1, a *C. elegans* β -catenin

homolog, which results in downregulation instead of upregulation of POP-1, a TCF/LEF transcription factor [20]. In contrast, noncanonical pathways act through cytoplasmic factors that ultimately regulate cytoskeletal components but can also act to regulate transcription independently of β -catenin [5]. Further understanding of the mechanisms by which Wnt signaling can regulate cell movements will be important for understanding morphogenesis during normal development and Wnt pathway function in tumor invasion and metastasis in human cancers [21].

We have used a candidate approach to begin to identify genes required for *C. elegans* gastrulation. We found that the pathways that specify the endodermal precursors also regulate ingression of these cells. We show that Wnt/Fz signaling has a role in gastrulation in addition to its function in regulating endodermal cell fate and cell-cycle timing. Although Wnt signaling functions in cell polarization in many contexts [22], Wnt signaling does not affect *C. elegans* gastrulation by establishing polarity in the ingressing cells or by affecting the rate of myosin accumulation at the apical cortex of these cells. Instead, we found that Wnt/Fz signaling functions in gastrulation by causing regulatory myosin light chain in the apical cortex of Ea and Ep to be phosphorylated at a contraction-activating serine residue. Our results forge new links among cell fate, cell signaling, and cell form and suggest a novel role for intercellular signaling by a Wnt protein—the regulation of morphogenetic movements through activation of myosin contractile activity.

Results

Endodermal-Fate-Specification Pathways Act Upstream of *C. elegans* Gastrulation

To determine whether endodermal fate specification is necessary for Ea and Ep morphogenetic behavior, we first determined whether *C. elegans* embryos defective in each of the endodermal-fate-specification genes also have gastrulation defects. The endodermal founder cells are specified by two intersecting pathways: A GATA-factor transcriptional cascade initially restricts mesoderm fate to the appropriate endoderm and mesoderm precursors, and the Wnt signaling pathway then acts to repress nuclear POP-1 and thus allow endoderm development in the progeny of the E cell [4]. The endoderm-specification pathways also affect cell-cycle timing in endodermal precursors. The Ea and Ep cells are the first cells in the *C. elegans* embryo to introduce a gap phase, a G₂ phase that results in the Ea and Ep cells dividing approximately 20 min later than MSa and MSp cells (20.2 ± 2.4 min SD; Figure 2B) [6]. This delay does not occur in endoderm-specification mutants [4]. We speculated that either endoderm specification or cell-cycle delay, or both, might be required for gastrulation. Gastrulation defects have been found in embryos defective in some of these genes, e.g., *mom-2* [23], *skn-1* [24], *end-1* [25], and *end-3* [26], but many of the relevant genetic backgrounds have not been analyzed for gastrulation movements.

Mutant embryos were imaged by 4D time-lapse video-microscopy and were subsequently analyzed for gastrulation movements. For the experiments described




Figure 2. *mom-2*/*Wnt*, *mom-5*/*Fz*, and *Dsh* Are Required for Gastrulation

(A) Gastrulation in wild-type embryos occurs with the ingress of Ea and Ep, and the embryos divide once they are completely surrounded by neighboring cells. Each panel is a single midsagittal optical section from a Nomarski time-lapse movie, with the minutes elapsed since Ea-Ep birth noted on the left side. All sections were examined to ensure that cells internalized in one section were internalized in three dimensions. E cells are pseudocolored green in all frames. *mom-2* and *mom-5* embryos are mutants, and the *Dsh* embryo shown is triple *dsh-1*;*dsh-2*;*mig-5*(RNAi). Arrowheads point to E lineage cells that failed to ingress. In this and all figures, embryos are oriented with anterior to the left and dorsal side up. Scale bar represents 5 μ m.

(B) There is no strict correlation between endoderm production and gastrulation or between Ea-Ep cell-cycle timing and gastrulation in *mom-2* or *mom-5*. Each dot represents one embryo (24 wild-type, 50 *mom-2*, and 34 *mom-5* embryos). The length of Ea-Ep delay, whether it gastrulated, and whether it produced endoderm are indicated.

in this paper, we defined successful gastrulation as the ingress of Ea and Ep to the point where they are completely surrounded by their neighbors before they divide (Figure 2A). We found that nearly all of the endoderm-specification mutants were also gastrulation defective, with rates of failure reaching as high as 100% in *mom-4*, *lit-1*, and *end-3* embryos (Table 1). The only endoderm-specification-pathway gene not required for gastrulation was *elt-2*, a downstream-acting

transcription factor whose expression is not detected until just after Ea-Ep ingress [27].

In an attempt to further identify potential links between patterning pathways and gastrulation, we determined whether ectopic endodermal cells in a mutant undergo ectopic gastrulation. PIE-1 is a CCCH zinc-finger protein that becomes segregated to posterior cells, where it specifies cell fates in part by repressing the function of the endomesoderm-specifying gene *skn-1*.

Table 1. Gastrulation Movements in Endoderm-Specification Mutants

Genotype	Homolog/Conserved Domains	Gastrulated/Total	Percentage Gastrulating	Allele Type ^a
Wild-type		17/17	100%	
<i>mom-1</i>	<i>porcupine</i>	2/18	11%	strong loss-of-function
<i>mom-2</i>	<i>wnt/wingless</i>	3/50	6%	deletion; genetic null
<i>mom-3</i>	<i>wntless</i>	3/7	43%	strong loss-of-function
<i>mom-4</i>	MEKK/TAK1	0/10	0%	strong loss-of-function
<i>mom-5</i>	<i>frizzled</i>	7/34	21%	TC1 transposon
<i>lit-1</i>	<i>nemo-like kinase</i>	0/10	0%	protein null
<i>pie-1</i>	CCCH zinc-finger protein	10/11 (Ea-Ep); 11/11 (P ₄ -D)*	91%	strong loss-of-function
<i>skn-1</i>	bZIP transcription factor	2/27	7%	strong loss-of-function
<i>end-1</i>	GATA factor	1/7	14%	deficiency; genetic null
<i>end-3</i>	GATA factor	0/12	0%	genetic null
<i>elt-2</i>	GATA factor	12/12	100%	TC1 transposon; genetic null

* Ea-Ep ingression is followed within the next 30 min by P₄-D ingression in *pie-1* embryos only. In wild-type, 0/17 embryos showed P₄-D ingression during this period.

^a References are as follows: *mom-1*, *mom-3*, and *mom-4* [23]; *mom-5* [33]; *lit-1* [55]; *pie-1* [55]; *skn-1* [24]; *end-1* [25]; *end-3* [25]; and *elt-2* [27]. See the Supplemental Experimental Procedures for allele designations.

Embryos from *pie-1* loss-of-function mothers produce ectopic endoderm from E's posterior neighbor, P₃, a cell that normally produces muscle and germline founder cells [25]. Time-lapse recordings of *pie-1* mutant embryos showed ectopic gastrulation: The two daughters of P₃ invariably ingressed soon after Ea-Ep (17.5 ± 3.3 min SD; Table 1). Because loss of function of endoderm-specifying genes interferes with gastrulation and loss of function of *pie-1*, a gene that prevents ectopic endoderm from forming, results in ectopic gastrulation, we conclude that the genes that specify endodermal cell fate in Ea and Ep function upstream of the cytoskeletal mechanisms that drive gastrulation. The most downstream player we found, the transcriptional activator END-1 [4], suggests that some molecular player(s) in gastrulation may be regulated transcriptionally, consistent with a known requirement for transcription in gastrulation [9].

Wnt/Frizzled Signaling Functions in Gastrulation as Well as Endoderm Specification

Next, we asked whether any of these endoderm-specification genes played a direct role in determining gastrulation behavior. Wnt ligands, Frizzled receptors, and Frizzled's downstream effectors function upstream of morphogenetic events in diverse organisms [15–18, 28]. We asked whether Wnt/Fz signaling functions during *C. elegans* gastrulation independently of its role in specifying endoderm by using rhabditiin granules as a terminal-differentiation marker for endoderm development (see the Supplemental Experimental Procedures available with this article online). If Wnt/Fz signaling affects gastrulation solely through its role in endoderm specification, then *mom-2*/Wnt or *mom-5*/Fz embryos that produce endoderm should gastrulate and those embryos that fail to make endoderm should fail to gastrulate. As expected, we found that all *mom-2* deletion allele embryos that failed to produce endoderm also failed to gastrulate (n = 33); however, we were surprised to find that most of the escapers—the endoderm-producing *mom-2* embryos—also failed to gastrulate (82%; n = 17) (Figure 2B). Similarly, most *mom-5* embryos that produced endoderm failed to gastrulate (85%; n = 34) (Figure 2B). We also examined the expression of

two endoderm-specific molecular markers, END-1, which is expressed during gastrulation, and ELT-2, which is expressed after gastrulation. *mom-5*(RNAi) embryos exhibited similar gastrulation defects to *mom-5*(*zu193*) embryos, and the END-1 and ELT-2 expression was comparable to that in wild-type cells in all embryos examined (n = 17 for END-1 and n = 9 for ELT-2; Movies S1 and S2 and Figure S1). Because Ea and Ep in *mom-5* embryos generally fail to gastrulate despite producing END-1 and producing ELT-2 and rhabditiin granules later, we propose that Wnt/Fz signaling has two functions: regulating endoderm specification and also, at least partially independently, regulating gastrulation.

The multidomain protein Dishevelled (Dsh) functions downstream of the Frizzled receptor in several systems [29]. *C. elegans* has three Dsh homologs: DSH-1, DSH-2, and MIG-5. Although null alleles of Dsh homologs affect endoderm specification, RNA interference (RNAi) of *dsh-1*, *dsh-2*, and *mig-5* individually or together has not [30]. This facilitated the determination of whether Dsh-mediated signaling is required for *C. elegans* gastrulation independently of a potential role in endoderm specification. We carried out RNAi of these three genes individually and in combination. With the exception of *dsh-1* RNAi, all of these treatments resulted in some embryos that failed to gastrulate despite producing endoderm, and RNAi of all three genes simultaneously produced a more penetrant gastrulation-defective phenotype than RNAi of any single gene alone (Figure 2 and Figure S2), suggesting that multiple Dsh proteins act redundantly to regulate gastrulation (we refer to these three proteins collectively as Dsh below). Our attempts to determine whether canonical or noncanonical signaling is involved in gastrulation downstream of Dsh have not yet resolved this issue. We conclude that MOM-2/Wnt, MOM-5/Fz, and multiple Dishevelled homologs function in the ingression of endodermal precursors in addition to their roles in endoderm specification. Because a *mom-2* null allele does not abolish gastrulation movements completely (Table 1), we conclude from these data that Wnt signaling acts partially redundantly as direct or indirect regulators of gastrulation with one or more additional pathways.

Wnt/Frizzled Signaling Affects Gastrulation Independently of Control of Ea-Ep Cell-Cycle Length

Previous work suggested that the G2 phase introduced in the Ea-Ep cell cycle near the time of gastrulation might be important for gastrulation: In *gad-1* (gastrulation defective) mutant embryos, endodermal cell fate is properly specified, but the Ea and Ep cells do not have a G2 phase nor do they ingress [31]. The division delay in Ea and Ep compared to their cousins MSa and MSP has been considered an aspect of endodermal cell fate because it is absent in many endoderm-deficient mutants [4]. We confirmed that the Ea-Ep division delay in *mom-2* embryos is generally shorter than that in wild-type embryos (Figure 2B). Very few *mom-2* embryos gastrulated (6%; n = 50), and the few embryos that did gastrulate had somewhat longer cell cycles than the average for *mom-2* embryos (8.5, 13, and 16.5 min compared with an average of 6.2 min). Similar results were seen in *mom-5* embryos and Dsh RNAi embryos (Figure 2B and Figure S2). These results demonstrate that embryos can gastrulate even when the Ea-Ep cell cycle is shorter than that in wild-type embryos, but they raised the possibility that more precocious cell division might prevent gastrulation movements, perhaps by reorganizing the actomyosin network for cell division in a way that precludes apical constriction from occurring at the same time.

We therefore conducted two types of experiments to test more directly whether Wnt signaling functions in gastrulation independently of its effect on cell-cycle timing. First, we lengthened the cell cycles of Ea and Ep in the absence of Wnt signaling. Ea and Ep cell cycles can be artificially lengthened by brief irradiation of Ea and Ep nuclei with a laser (referred to here as laser-delay; see Supplemental Experimental Procedures). This treatment did not interfere with ingression in wild-type embryos (9/9 cases; Figure 3A). In *gad-1* embryos, laser-delaying Ea and Ep cell division rescued ingression in approximately half of the cases (9/19 cases; Figure 3A), and control laser-delay of cell pairs in AB or P₂ lineages in *gad-1* embryos did not cause ingression (data not shown). The rescue of ingression by laser-delay of Ea and Ep suggests that *gad-1* functions in gastrulation only indirectly, through regulating Ea and Ep division timing, and also indicates that laser-delay can be used to rescue ingression in a mutant that would otherwise have premature Ea-Ep division.

In contrast to the *gad-1* experiments, irradiation of Ea and Ep nuclei to delay Ea and Ep cell division completely failed to rescue gastrulation movements in *mom-2* embryos (0/14 cases; Figure 3A). Because neither long Ea-Ep cell cycles in untreated *mom-5* embryos nor artificially extended Ea-Ep cell cycles in *mom-2* embryos are sufficient for gastrulation to occur, and yet many such embryos would normally produce endoderm (Figure 3 legend), we propose that endoderm specification and long Ea-Ep cell cycles may not be sufficient for gastrulation to occur in the absence of this Wnt signaling pathway, and that Wnt signaling has additional functions in gastrulation.

Second, if Wnt signaling acts once to specify endoderm and cell-cycle timing and again later during gastrulation, then manipulating the presence of a Wnt signal after induction of endodermal fate and cell-cycle timing




Figure 3. *mom-2* Is Required for Gastrulation in Addition to its Effect on Ea-Ep Cell-Cycle Timing

(A) Laser-mediated delay of Ea-Ep cell division does not prevent gastrulation in wild-type embryos (9/9 cell pairs ingressed), and it can sometimes rescue gastrulation in *gad-1* embryos (9/19 cell pairs), but it cannot rescue gastrulation in *mom-2* embryos (0/14 cell pairs, significantly lower than the proportion of untreated *mom-2* embryos that produced endoderm in Figure 2, chi-square test, $p < 0.05$). A schematic of MSa-Ep and Ea-Ep (green) cell positions is shown above cell-division patterns. Lightning bolts represent approximate time of irradiation.

(B) In vitro cell-recombination experiments suggest that MOM-2 functions in gastrulation after endoderm specification has occurred. Three experiments are diagrammed at bottom. In these experiments, wild-type P₂ cells recombined with *mom-2* EMS cells always rescued endoderm and rescued normal cell-cycle division timing in Ea-Ep cells. The top of the panel shows quarter-pie graphs depicting the extent of movement by MSxx relative to Ea-Ep during the 30 min after MSxx birth in each experiment. Each short line represents the degree of movement in one recombination experiment. The black area encompasses 50% of the data, and the gray area encompasses a further 25% of the data. Results are quantified in the bar graph below, in which the proportions that gastrulated are shown (more than 8° of MSxx movement was required to score movement as gastrulation, based on previous results [7]). Wild-type P₂ cells more successfully rescued movement. The asterisk indicates that the results from this manipulation was statistically different from each of the other two manipulations with chi-square tests, at $p < 0.05$.

should determine whether gastrulation occurs. At the four-cell stage, P₂ induces endoderm in the neighboring EMS cell via Wnt signaling [23, 32, 33]. The timing of this induction is well characterized and can be manipulated in an in vitro blastomere culture system [34]. We used these cell manipulations to present a Wnt signaling cell

to specify E cell fate, and we then replaced it with either Wnt-minus or Wnt-plus signaling cells to look for specific effects on gastrulation. First, a number of control experiments were performed. When we separated wild-type P₂ from wild-type EMS cells and recombined these cells with wild-type partners that were at the same developmental time point, most recombinants exhibited gastrulation movements (75%, n = 16; [Figure 3B](#)). In contrast, *mom-2* P₂ and *mom-2* EMS negative control recombinations resulted in little or no movement ([Figure S3](#)), as seen previously in related experiments with *mom-2* [7]. We then confirmed that gastrulation defects could be rescued by recombining *mom-2* EMS cells with wild-type P₂ cells (70%, n = 20; [Figure 3B](#)). Next, we asked whether a Wnt signaling cell during endoderm specification was sufficient for gastrulation movements by replacing a Wnt-plus P₂ cell with Wnt-minus cells after endoderm induction. We found that after endoderm induction had occurred, a Wnt-plus P₂ cell is able to rescue gastrulation movements (70%, n = 20) significantly more effectively than a Wnt-minus signaling cell (36%, n = 14; [Figure 3B](#)), suggesting that the P₂ cell or its descendants, or both, are likely to be a source of Wnt signaling for gastrulation.

Together, these results suggest that although Wnt signaling can affect gastrulation indirectly by regulating endodermal cell fate and division timing, Wnt signaling also has a second role during gastrulation. This could be a second, independent Wnt-Fz interaction, or possibly a higher threshold response to the interaction that establishes endoderm cell fate. These results prompted us to look for more direct cellular effects of Wnt/Fz signaling during gastrulation by examining the effect of Wnt/Fz signaling on the polarization and cytoskeletal motility of ingressing cells. Below, we show that in the absence of Wnt signaling, some cell biological events implicated in ingression occur normally in Ea and Ep, but others do not.

Ea and Ep Apicobasal Polarization Proceeds Independently of Wnt/Frizzled Signaling

Because Wnt and Fz-dependent signaling are known to affect cell polarity in several systems [35], we examined whether Ea-Ep apicobasal polarity was disrupted in the absence of Wnt/Fz signaling. During gastrulation, the PAR proteins are localized in apicobasally polarized patterns and are required for apical myosin enrichment and efficient ingression movements [8, 10]. We examined wild-type embryos expressing PAR-2::GFP and compared them to PAR-2::GFP embryos from mothers that were either fed with a bacterial strain expressing *mom-2* dsRNA or injected with *mom-5* dsRNA. PAR-2::GFP;*mom-2*(RNAi) embryos and PAR-2::GFP;*mom-5*(RNAi) embryos exhibited similar gastrulation defects to *mom-2*(*or309*) and *mom-5*(*zu193*) embryos respectively (see [Supplemental Experimental Procedures](#)). We confirmed that GFP is detected in PAR-2::GFP embryos in a basolateral pattern in Ea-Ep (11/11 embryos; [Figure 4A](#)). We found that PAR-2::GFP;*mom-2*(RNAi) embryos also display PAR-2::GFP in a basolateral pattern (16/16 embryos; [Figure 4B](#)) as do PAR-2::GFP;*mom-5*(RNAi) embryos (8/8 embryos; [Figure 4C](#)), indicating that Wnt-Fz signaling does not regulate PAR-2 basolateral distribution at this stage.




Figure 4. GFP::PAR-2 Distribution Shows that Apicobasal Polarity Is Established Normally in the Absence of *mom-2*/Wnt or *mom-5*/Fz. GFP::PAR-2 is enriched basolaterally in Ea and Ep in wild-type (A), *mom-2* (B), and *mom-5* (C) embryos. Ea-Ep are labeled with asterisks. Arrows mark the basal sides of Ea-Ep cells, where GFP::PAR-2 is enriched, and arrowheads mark the apical sides. P₄, on the posterior side of the embryo, exhibits nonlocalized GFP::PAR-2 because the GFP construct is driven by a PIE-1 promoter that is most active in the P lineage. Confocal images are shown; scale bars represent 5 μ m.

A second step in cell polarization occurs when NMY-2, a nonmuscle myosin II heavy chain, accumulates at the apical surfaces of Ea and Ep, where it contributes to gastrulation movements [7, 8]. It is possible that Wnt/Fz signaling may effect gastrulation by acting in parallel with PAR proteins to enrich NMY-2 apically. The majority of cortical NMY-2 accumulation in Ea and Ep occurs during the G2 phase of the Ea and Ep cells [8]. We used *mom-5* mutants as a source of Wnt signaling-deficient embryos because Ea-Ep division timing in *mom-5* mutant embryos more closely resembles that in wild-type embryos than it does in *mom-2* mutants ([Figure 2B](#)). We found that NMY-2::GFP accumulated in the apical cortex of Ea and Ep in *mom-5* embryos as much as in wild-type embryos and did not accumulate apically in other cells at this stage ([Figure 5A](#)). Quantification of cortical to cytoplasmic NMY-2::GFP ratios confirmed this: Wild-type and *mom-5* Ea-Ep cells accumulated NMY-2::GFP at similar rates, and both accumulated significantly more NMY-2::GFP than did cells not of E lineage in either background ([Figure 5B](#)). It is therefore unlikely that Wnt/Fz signaling affects gastrulation by regulating the accumulation of NMY-2. These results




Figure 5. Apical NMY-2 Accumulation Occurs in Both Wild-Type and *mom-5*/Fz Ea and Ep Cells and at Similar Rates

(A) Confocal images from recordings of wild-type and *mom-5* embryos expressing NMY-2::GFP. Asterisks mark Ea and Ep cells. d represents distance, t represents time, where t_i is the initial time point analyzed and t_f is the final time point analyzed 25 min later. Boxes around the Ea and Ep cortex represent the areas used to make each kymograph at right, in which the maximum pixel value along each d position in each box is shown over 25 min. In the kymographs, the cortex in wild-type can be seen accumulating NMY-2::GFP and moving toward the interior of the embryo, and the cortex in *mom-5* can be seen accumulating NMY-2::GFP but moving only erratically. Similar areas were analyzed on three other quadrants of the embryo.

(B) Graph of cortical to cytoplasmic NMY-2::GFP level over time. Five wild-type embryos, and four *mom-5* embryos that failed to gastrulate, were analyzed. NMY-2::GFP levels were quantified by calculation of the ratio of cortical to cytoplasmic pixel level above background. A ratio of one therefore indicates that the cortex is at the same intensity as the cytoplasm. By 25 min after P₄ birth, E cell cortices have enriched significantly more NMY-2::GFP than non-E cells, in both backgrounds ($p < 0.05$). The wild-type Ea-Ep GFP ratio is not statistically distinguishable from the *mom-5* Ea-Ep GFP ratio at either time point. Error bars indicate ± 1 SD.

suggest that Wnt/Fz signaling does not appear to affect gastrulation through regulation of apicobasal polarity in the ingressing cells.

Frizzled Signaling Functions Upstream of Apical Constriction

Because myosin II heavy chain becomes enriched on the apical side of the ingressing cells at a normal rate but ingression does not occur in Wnt-pathway mutants,




Figure 6. Apical Domains of Ea-Ep Fail to Constrict in *mom-5* Mutant Embryos

Lengths of the apical domains of Ea and Ep over time are shown for (A) 13 wild-type and (B) six *mom-5* mutant embryos that produced endoderm but failed to gastrulate. Error bars are 95% confidence bars.

we speculated that Wnt signaling might instead regulate contraction of the apical actomyosin network. To test this, we first measured the rate of apical constriction during gastrulation in wild-type and *mom-5* (RNAi) embryos. We generated 4D movies of gastrulating embryos and marked the sites of contact between Ea and Ep and the surrounding cells at their apical surfaces. We then calculated the distances between these sites of cell contact. We considered Ea-Ep apical-surface measurements for only the first 10 min after MSa-MSp division so that the premature Ea-Ep division that occurs in some *mom-5* embryos could not affect the results. During this time in wild-type embryos, the length of the apical domain decreased at a rate of 7.1 ± 1.3 nm/s (Figure 6). In contrast, the length of the apical domain of the Ea and Ep cells in *mom-5* embryos that produced endoderm but failed to internalize Ea and Ep decreased significantly more slowly, at a rate of 2.3 ± 3.6 nm/s ($p < 0.05$). These data suggest that Wnt signaling regulates constriction of the apical domains of the Ea and Ep cells.

Myosin II Becomes Activated in the Apical Cortex of the Ea and Ep Cells Near the Time of Ingression

The failure of apical constriction in the absence of Wnt signaling suggested that apical myosin II might normally become activated to contract at the time of gastrulation and that Wnt signaling might function in this activation. Myosin II has two heavy chains and two essential light chains and two regulatory light chains, and phosphorylation of the two regulatory light chains (referred to here as rMLC) at serine 19 is required for formation of active myosin filaments that can drive contraction in various systems [36]. We used an antibody that recognizes this phosphoepitope, which we refer to as p-rMLC, to determine the localization of activated myosin before and during gastrulation. We found first that commonly




Figure 7. Phosphorylated Regulatory Myosin Light Chain Accumulates Apically in the Endoderm Precursors Near the Time that Gastrulation Occurs

Gastrulation movements begin near the 26-to 28-cell stage in most embryos. Ea-Ep cells are indicated by asterisks, and apical-domain membranes are labeled with arrowheads. p-rMLC is shown in green, F-actin is shown in red, and DAPI staining is shown in blue.

used methods for immunostaining *C. elegans* embryos with methanol failed to preserve the p-rMLC signal, but the signal could be detected reliably with a formaldehyde fix designed to best preserve actin filaments [37] (see Supplemental Experimental Procedures). One-cell embryos stained in accordance with this protocol showed cortical staining consistent with published results [38, 39], and cortical staining was abolished by RNAi to the rMLC-encoding gene *mlc-4* (16/16 embryos; Figures S4 and S5). Furthermore, the cortical staining colocalized with NMY-2, the myosin heavy chain (Figure S6), suggesting that the antibody we used can recognize p-rMLC at the cell cortex. During gastrulation, we detected a striking enrichment of p-rMLC at the apical surfaces of Ea and Ep (Figure 7). p-rMLC staining appears enriched apically in the ingressing Ea and Ep cells specifically starting at the 26-cell stage (after MSA-MSp division) and until the Ea and Ep cells are fully ingressed, and it is not enriched apically in any other nondividing cells during these stages. We conclude that apically localized regulatory myosin light chain is activated in Ea and Ep near the time of ingression, by an as yet unidentified kinase.

Frizzled Signaling Activates Myosin II at the Ea and Ep Apical Cortex

Next, we asked whether Wnt-Fz signaling functions upstream of rMLC phosphorylation by immunostaining *mom-5* embryos with anti-p-rMLC. In light of the fact that Ea and Ep cells can divide earlier in *mom-5* embryos than in wild-type, we were careful to only compare wild-type and *mom-5* embryos of similar stages (Figure 8 and Supplemental Experimental Procedures). In contrast with wild-type embryos, *mom-5* embryos exhibited markedly lower levels of p-rMLC staining at the apical surface of Ea and Ep throughout all stages of gastrulation movements (Figure 8). The apical surfaces of Ea and Ep in *mom-5* embryos stained in few cases, and staining was weak in these cases (2/11 embryos), whereas apical staining in wild-type embryos is more robust (10/20 embryos showing strong staining, as shown in the Figure 8, and 6/20 embryos showing





Figure 8. Regulatory Myosin Light-Chain Phosphorylation Depends on MOM-5/Fz

Wild-type embryos show apical accumulation of phosphorylated rMLC as gastrulation occurs, particularly in 26- and 28-cell stage embryos. In contrast, *mom-5* embryos show reduced accumulation of phosphorylated rMLC compared to wild-type embryos of the same stages. Ea-Ep cells are indicated by asterisks. p-rMLC is shown in green, F-actin is shown in red, and DAPI is shown in blue. In the bottom panels, Ea-Ep apical membranes are expanded to show p-rMLC staining. Apical membranes are denoted by arrowheads.

weak staining equivalent to the *mom-5* embryos). We conclude that in the absence of Wnt signaling, there is a significant reduction in the level of phosphorylated regulatory myosin light chain on the apical sides of the Ea and Ep cells.

Discussion

The study of morphogenesis is one of the key areas where cell and developmental biology meet [1, 40]. Although embryonic patterning mechanisms can play crucial roles in determining cytoskeletal behaviors, there are few examples where the coupling between embryonic patterning and cytoskeletal behaviors is well understood [41]. We have found that the well-studied pathways that specify endoderm in *C. elegans* act upstream of mechanisms that internalize the endodermal precursors. We have demonstrated that Wnt-Fz signaling regulates *C. elegans* gastrulation in addition to specifying endodermal cell fate and cell-cycle timing. The mechanism by which Wnt-Fz signaling acts is not through the generation of the polarized distribution of PAR proteins



or apical-myosin accumulation, suggesting that Wnt-Fz signaling might directly affect apical constriction downstream of myosin accumulation. In support of this, we found an enrichment of an activated form of myosin at the apical cortex of wild-type Ea and Ep during gastrulation, and we found that this activation of myosin is dependent on signaling through Fz.

Our results rule out the simple possibility that Wnt signaling affects *C. elegans* gastrulation solely through its well-documented role in endoderm specification because we found that Ea and Ep often produce endoderm by multiple measures, but fail to gastrulate, in the absence of Wnt signaling. However, it remains possible that rMLC phosphorylation could be a higher threshold response to Wnt-Fz signaling at the four-cell stage. Alternatively, rMLC phosphorylation could depend on an independent, later Wnt-Fz interaction. Signaling to rMLC downstream of Dsh in either way could occur by a transcriptional mechanism or by signaling that is more direct to rMLC. It will be of interest to identify more members of this signaling pathway and to determine when they function so that these questions can be answered.

Our results, together with previous results from us and others [7, 9, 10] outline a molecular and mechanical model for *C. elegans* gastrulation (Figure 9). The apico-basal polarity of all or most cells is determined by the positions of cell-cell contacts and is reflected in the localization of PAR proteins. The pathways that specify endodermal fate in *C. elegans*, involving many of the genes listed in Table 1, determine which cells will enrich myosin heavy chain apically in response to PAR-protein localization. Having myosin enriched at the apical, contact-free surfaces likely primes these cells for internalization. Wnt-Fz signaling leads to phosphorylation of a conserved residue on rMLC, and this phosphorylation can result in the formation of active myosin filaments. The contraction of the apical actomyosin machinery then shrinks the contact-free apical areas, pulls neighboring cells under the Ea and Ep cells, and results in the ingression of Ea and Ep into the center of the embryo. In this way, ingression appears to depend on the combinatorial information from cell-fate specification, apico-basal polarity, and reception of a cell signal that results in contraction of the actomyosin network on a specific side of specific cells.

The role of cell-fate-specification mechanisms and Wnt signaling in activating ingression of specific cells in *C. elegans* gastrulation has both striking parallels and critical differences with *Drosophila* gastrulation. In *Drosophila*, the mesoderm-specification protein Twist activates apical secretion of a different intercellular signaling protein, Folded-gastrulation (Fog) [41, 42]. Fog acts through a presumed but unidentified cell-surface receptor and a G α protein, Concertina, to cause myosin localization to the apical surface of cells. Loss of the Fog downstream targets, DRhoGEF or the Rho kinase Drok, causes much more severe defects in ventral furrow formation than does loss of Concertina, suggesting that Fog must act redundantly with another, unidentified pathway to drive apical constriction [41]. Myosin motor activity is required for its apical localization in *Drosophila* gastrulation [41]. Whether Fog signaling regulates myosin activity by rMLC phosphorylation in *Drosophila* gastrulation like the Wnt pathway does in *C. elegans* has not been examined, but Rho kinase is known to cause phosphorylation of rMLC or myosin activation in other settings in *Drosophila* [43–45] and other organisms [36].

In vertebrates, the actin-binding protein Shroom functions to localize both actin filaments and myosin and can cause apical constriction when expressed in MDCK cells and during normal neural-tube closure [46–48]. Shroom functions in apical constriction by restricting myosin localization, and it is not known to affect myosin activity. Shroom acts through a small GTPase like Fog, but does so through a different GTPase, Rap1. Fog, Shroom, and Wnt-dependent control of apical constriction appear to work independently rather than as parts of a single pathway because Fog and Shroom do not have homologs in the *C. elegans* genome, and removal of the function of the Rap1 homolog in *C. elegans* has no effect on gastrulation (T. Grana and J.H., unpublished data). Also, Shroom is required for apical constriction of only some cells in the organisms where it functions; for example, *Xenopus* bottle cells do not require Shroom during gastrulation [48]. These data suggest that apical constriction is regulated during animal development by multiple, independent mechanisms that can affect either myosin distribution or activity, or both.

Although ours is the first report of intercellular Wnt signaling regulating morphogenesis through rMLC phosphorylation, there is precedent for Frizzled acting

in a similar but cell-autonomous fashion in *Drosophila*, suggesting that this might be an ancient mechanism of morphogenetic regulation. Frizzled can affect ommatidial polarity and actin bundle number in *Drosophila* planar-cell polarity (PCP) signaling pathways via Dsh, RhoA, Drok, and rMLC phosphorylation [45]. This pathway suggests some molecular players that might act between Dsh and rMLC phosphorylation in *C. elegans* gastrulation. The *Drosophila* PCP pathways differ from what we have outlined in this report in that Frizzled is probably not responding to intercellular Wnt signals in PCP signaling pathways [49, 50] and in that PCP signaling is not known to drive apical constriction. The mechanism by which myosin activation regulates ommatidial polarity and actin bundle number in PCP pathways is unknown.

In zebrafish and frog embryos, PCP signaling is required during gastrulation for convergence and extension movements [17, 18]. Vertebrate PCP genes are known to regulate the activity of the cytoskeletal modifiers Rho and Rac [51–53], but the direct effect of PCP signaling on the cytoskeleton has not yet been analyzed in detail. Both the vertebrate PCP pathway and *C. elegans* Wnt/Fz signaling result in modification of cell shape and cell behavior during gastrulation. These pathways differ in that vertebrate PCP signaling affects cell polarity [17, 18], whereas cell polarity is undisturbed in *mom-2/Wnt* and *mom-5/Fz* mutant embryos (this report) and RNAi experiments targeting *C. elegans* homologs of the PCP genes *fat* and *flamingo* (D.M., unpublished) and *van gogh/strabismus* (T.W. and J.H., unpublished data) have not resulted in any gastrulation defects, suggesting that *C. elegans* Wnt/Fz signaling in gastrulation through conventional PCP signaling is unlikely.

Why would myosin activity in *C. elegans* gastrulation be regulated by a cell-cell signal? It seems plausible that constitutively activating myosin could achieve the same goal because myosin is enriched in the apical cortex of the Ea and Ep cells, and hence, activation of myosin throughout these cells might cause an imbalance of forces that could drive apical constriction reliably. One possible explanation is that signaling ensures that apical constriction occurs at the right time. For example, signaling could ensure that myosin activation occurs at a time when the actomyosin network is apically enriched and is not being used for cell division or at a time when adhesion to neighboring cells is sufficiently strong for constriction to pull neighboring cells under the Ea and Ep cells. Signaling might also contribute spatial specificity. For example, it is possible that there are stages in development when several cells have the potential to ingress but only the appropriate ones do so because they contact a Wnt signaling cell.

Others have speculated that morphogenesis may more often depend on redundant pathways than embryonic patterning does [2]. Very few backgrounds that we examined prevented gastrulation in all embryos, suggesting that *C. elegans* gastrulation is regulated by multiple, partially redundant mechanisms. Morphogenesis depends on diverse mechanisms that are of interest in the field of cell and developmental biology, including spatial and temporal gene regulation, cell signaling, cell polarization, cell adhesion, and cytoskeletal dynamics. It will be of interest to explore how such

mechanisms work together in morphogenesis in *C. elegans* gastrulation.

Conclusions

We have used a combination of genetics, cell manipulations, and in vivo imaging to investigate the regulation of the cytoskeleton by cell-fate-specification genes and by intercellular signaling pathways during the morphogenetic movements of *C. elegans* gastrulation. We have shown that the pathways required for endoderm specification, including a Wnt/Frizzled pathway, are required for gastrulation to occur. Furthermore, we have shown that this Wnt/Frizzled pathway functions in gastrulation in addition to specifying endodermal cell fate. In the absence of Frizzled, apical constriction of the endoderm precursors fails to occur. Additionally, embryos lacking Frizzled show reduced levels of phosphorylated myosin in the apical domains of the endodermal precursors compared to wild-type embryos. Because this phosphorylation of myosin is likely to drive actomyosin contraction, and thus the ingression in these cells at this site, we hypothesize that this failure to phosphorylate myosin underpins the gastrulation defect in Wnt/Frizzled signaling defective embryos. Thus, we have demonstrated a novel role for Wnt signaling during morphogenesis through understanding its modulation of the cytoskeleton and how this impacts upon cell movements.

Supplemental Data

Supplemental Data include six figures and two movies and can be found with this article online at <http://www.current-biology.com/cgi/content/full/16/20/1986/DC1/>.

Acknowledgments

We thank Mark Peifer, Steve Rogers, John Wallingford and members of the Goldstein lab for useful discussions and encouragement and Jeremy Nance, Ed Munro, Jim Priess, Morris Maduro, and the Caenorhabditis Genetics Center for worm strains. The Caenorhabditis Genetics Center is supported by the National Institutes of Health National Center for Research Resources. This work was supported by National Science Foundation grant IOB0518081 to J.H. and National Institutes of Health R01-GM68966 to B.G.

Received: February 21, 2006

Revised: August 21, 2006

Accepted: August 25, 2006

Published: October 23, 2006

References

1. Dawes-Hoang, R.E., and Wieschaus, E.F. (2001). Cell and developmental biology—a shared past, an intertwined future. *Dev. Cell* 1, 27–36.
2. Wieschaus, E. (1995). From molecular patterns to morphogenesis: The lessons from *Drosophila*. In *Nobel Lectures, Physiology or Medicine 1991–1995*, N. Ringertz, ed. (Singapore: World Scientific Publishing Co.)
3. Sulston, J.E., Schierenberg, E., White, J.G., and Thomson, J.N. (1983). The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev. Biol.* 100, 64–119.
4. Maduro, M.F., and Rothman, J.H. (2002). Making worm guts: The gene regulatory network of the *Caenorhabditis elegans* endoderm. *Dev. Biol.* 246, 68–85.
5. Thorpe, C.J., Schlesinger, A., and Bowerman, B. (2000). Wnt signaling in *Caenorhabditis elegans*: Regulating repressors and polarizing the cytoskeleton. *Trends Cell Biol.* 10, 10–17.

6. Edgar, L.G., and McGhee, J.D. (1988). DNA synthesis and the control of embryonic gene expression in *C. elegans*. *Cell* 53, 589–599.
7. Lee, J.Y., and Goldstein, B. (2003). Mechanisms of cell positioning during *C. elegans* gastrulation. *Development* 130, 307–320.
8. Nance, J., and Priess, J.R. (2002). Cell polarity and gastrulation in *C. elegans*. *Development* 129, 387–397.
9. Nance, J., Lee, J.-Y., and Goldstein, B. Gastrulation in *C. elegans*, in *WormBook* (The *C. elegans* Research Community, <http://www.wormbook.org>). 2005.
10. Nance, J., Munro, E.M., and Priess, J.R. (2003). *C. elegans* PAR-3 and PAR-6 are required for apicobasal asymmetries associated with cell adhesion and gastrulation. *Development* 130, 5339–5350.
11. Kemphues, K. (2000). PARsing embryonic polarity. *Cell* 101, 345–348.
12. Wodarz, A. (2002). Establishing cell polarity in development. *Nat. Cell Biol.* 4, E39–E44.
13. Logan, C.Y., and Nusse, R. (2004). The Wnt signaling pathway in development and disease. *Annu. Rev. Cell Dev. Biol.* 20, 781–810.
14. Peifer, M., Rauskolb, C., Williams, M., Riggleman, B., and Wieschaus, E. (1991). The segment polarity gene armadillo interacts with the wingless signaling pathway in both embryonic and adult pattern formation. *Development* 111, 1029–1043.
15. Wieschaus, E., and Riggleman, R. (1987). Autonomous requirements for the segment polarity gene armadillo during *Drosophila* embryogenesis. *Cell* 49, 177–184.
16. Gong, Y., Mo, C., and Fraser, S.E. (2004). Planar cell polarity signalling controls cell division orientation during zebrafish gastrulation. *Nature* 430, 689–693.
17. Heisenberg, C.P., Tada, M., Rauch, G.J., Saude, L., Concha, M.L., Geisler, R., Stemple, D.L., Smith, J.C., and Wilson, S.W. (2000). Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. *Nature* 405, 76–81.
18. Wallingford, J.B., Rowing, B.A., Vogeli, K.M., Rothbacher, U., Fraser, S.E., and Harland, R.M. (2000). Dishevelled controls cell polarity during *Xenopus* gastrulation. *Nature* 405, 81–85.
19. Wallingford, J.B., and Harland, R.M. (2002). Neural tube closure requires Dishevelled-dependent convergent extension of the midline. *Development* 129, 5815–5825.
20. Herman, M.A., and Wu, M. (2004). Noncanonical Wnt signaling pathways in *C. elegans* converge on POP-1/TCF and control cell polarity. *Front. Biosci.* 9, 1530–1539.
21. Vincan, E. (2004). Frizzled/WNT signalling: The insidious promoter of tumour growth and progression. *Front. Biosci.* 9, 1023–1034.
22. Sancho, E., Batlle, E., and Clevers, H. (2004). Signaling pathways in intestinal development and cancer. *Annu. Rev. Cell Dev. Biol.* 20, 695–723.
23. Thorpe, C.J., Schlesinger, A., Carter, J.C., and Bowerman, B. (1997). Wnt signaling polarizes an early *C. elegans* blastomere to distinguish endoderm from mesoderm. *Cell* 90, 695–705.
24. Bowerman, B., Eaton, B.A., and Priess, J.R. (1992). *skn-1*, a maternally expressed gene required to specify the fate of ventral blastomeres in the early *C. elegans* embryo. *Cell* 68, 1061–1075.
25. Zhu, J., Hill, R.J., Heid, P.J., Fukuyama, M., Sugimoto, A., Priess, J.R., and Rothman, J.H. (1997). *end-1* encodes an apparent GATA factor that specifies the endoderm precursor in *Caenorhabditis elegans* embryos. *Genes Dev.* 11, 2883–2896.
26. Maduro, M.F., Hill, R.J., Heid, P.J., Newman-Smith, E.D., Zhu, J., Priess, J.R., and Rothman, J.H. (2005). Genetic redundancy in endoderm specification within the genus *Caenorhabditis*. *Dev. Biol.* 284, 509–522.
27. Fukushige, T., Hawkins, M.G., and McGhee, J.D. (1998). The GATA-factor *elt-2* is essential for formation of the *Caenorhabditis elegans* intestine. *Dev. Biol.* 198, 286–302.
28. McEwen, D.G., Cox, R.T., and Peifer, M. (2000). The canonical Wg and JNK signaling cascades collaborate to promote both dorsal closure and ventral patterning. *Development* 127, 3607–3617.
29. Wallingford, J.B., and Habas, R. (2005). The developmental biology of Dishevelled: An enigmatic protein governing cell fate and cell polarity. *Development* 132, 4421–4436.
30. Walston, T., et al. (2004). Multiple Wnt signaling pathways converge to orient the mitotic spindle in early *C. elegans* embryos. *Dev. Cell* 7, 831–841.
31. Knight, J.K., and Wood, W.B. (1998). Gastrulation initiation in *Caenorhabditis elegans* requires the function of *gad-1*, which encodes a protein with WD repeats. *Dev. Biol.* 198, 253–265.
32. Goldstein, B. (1992). Induction of gut in *Caenorhabditis elegans* embryos. *Nature* 357, 255–257.
33. Rocheleau, C.E., Downs, W.D., Lin, R., Wittmann, C., Bei, Y., Cha, Y.H., Ali, M., Priess, J.R., and Mello, C.C. (1997). Wnt signaling and an APC-related gene specify endoderm in early *C. elegans* embryos. *Cell* 90, 707–716.
34. Goldstein, B. (1993). Establishment of gut fate in the E lineage of *C. elegans*: The roles of lineage-dependent mechanisms and cell interactions. *Development* 118, 1267–1277.
35. Axelrod, J.D., and McNeill, H. (2002). Coupling planar cell polarity signaling to morphogenesis. *ScientificWorldJournal* 2, 434–454.
36. Somlyo, A.P., and Somlyo, A.V. (2003). Ca²⁺ sensitivity of smooth muscle and nonmuscle myosin II: Modulated by G proteins, kinases, and myosin phosphatase. *Physiol. Rev.* 83, 1325–1358.
37. Costa, M., Draper, B.W., and Priess, J.R. (1997). The role of actin filaments in patterning the *Caenorhabditis elegans* cuticle. *Dev. Biol.* 184, 373–384.
38. Jenkins, N., Saam, J.R., and Mango, S.E. (2006). CYK-4/GAP provides a localized cue to initiate anteroposterior polarity upon fertilization. *Science* 313, 1298–1301.
39. Piekny, A.J., and Mains, P.E. (2002). Rho-binding kinase (LET-502) and myosin phosphatase (MEL-11) regulate cytokinesis in the early *Caenorhabditis elegans* embryo. *J. Cell Sci.* 115, 2271–2282.
40. Keller, R. (2002). Shaping the vertebrate body plan by polarized embryonic cell movements. *Science* 298, 1950–1954.
41. Dawes-Hoang, R.E., Parmar, K.M., Christiansen, A.E., Phelps, C.B., Brand, A.H., and Wieschaus, E.F. (2005). Folded gastrulation, cell shape change and the control of myosin localization. *Development* 132, 4165–4178.
42. Costa, M., Wilson, E.T., and Wieschaus, E. (1994). A putative cell signal encoded by the folded gastrulation gene coordinates cell shape changes during *Drosophila* gastrulation. *Cell* 76, 1075–1089.
43. Mizuno, T., Tsutsui, K., and Nishida, Y. (2002). *Drosophila* myosin phosphatase and its role in dorsal closure. *Development* 129, 1215–1223.
44. Rogers, S.L., Wiedemann, U., Hacker, U., Turck, C., and Vale, R.D. (2004). *Drosophila* RhoGEF2 associates with microtubule plus ends in an EB1-dependent manner. *Curr. Biol.* 14, 1827–1833.
45. Winter, C.G., Wang, B., Ballew, A., Royou, A., Kares, R., Axelrod, J.D., and Luo, L. (2001). *Drosophila* Rho-associated kinase (Drok) links Frizzled-mediated planar cell polarity signaling to the actin cytoskeleton. *Cell* 105, 81–91.
46. Hildebrand, J.D., and Soriano, P. (1999). Shroom, a PDZ domain-containing actin-binding protein, is required for neural tube morphogenesis in mice. *Cell* 99, 485–497.
47. Hildebrand, J.D. (2005). Shroom regulates epithelial cell shape via the apical positioning of an actomyosin network. *J. Cell Sci.* 118, 5191–5203.
48. Haigo, S.L., Hildebrand, J.D., Harland, R.M., and Wallingford, J.B. (2003). Shroom induces apical constriction and is required for hinge point formation during neural tube closure. *Curr. Biol.* 13, 2125–2137.
49. Strutt, D. (2003). Frizzled signalling and cell polarisation in *Drosophila* and vertebrates. *Development* 130, 4501–4513.
50. Lawrence, P.A., Casal, J., and Struhl, G. (2002). Towards a model of the organisation of planar polarity and pattern in the *Drosophila* abdomen. *Development* 129, 2749–2760.
51. Habas, R., Kato, Y., and He, X. (2001). Wnt/Frizzled activation of Rho regulates vertebrate gastrulation and requires a novel Formin homology protein Daam1. *Cell* 107, 843–854.
52. Habas, R., Dawid, I.B., and He, X. (2003). Coactivation of Rac and Rho by Wnt/Frizzled signaling is required for vertebrate gastrulation. *Genes Dev.* 17, 295–309.

53. Marlow, F., Topczewski, J., Sepich, D., and Solnica-Krezel, L. (2002). Zebrafish Rho kinase 2 acts downstream of Wnt11 to mediate cell polarity and effective convergence and extension movements. *Curr. Biol.* 12, 876–884.
54. Park, F.D., Tenlen, J.R., and Priess, J.R. (2004). *C. elegans* MOM-5/frizzled functions in MOM-2/Wnt-independent cell polarity and is localized asymmetrically prior to cell division. *Curr. Biol.* 14, 2252–2258.
55. Meneghini, M.D., Ishitani, T., Carter, J.C., Hisamoto, N., Nino-miya-Tsuji, J., Thorpe, C.J., Hamill, D.R., Matsumoto, K., and Bowerman, B. (1999). MAP kinase and Wnt pathways converge to downregulate an HMG-domain repressor in *Caenorhabditis elegans*. *Nature* 399, 793–797.